



NORTHWEST EYE SURGEONS

Tel. 800.826.4631

www.nweyes.com



WHATCOM EYE SURGEONS

A Division of Northwest Eye Surgeons of Seattle

www.whatcomeyes.com

Vision Correction 2: Multifocal IOL Post-operative Care

Best results with multifocal vision are achieved after both eyes are implanted.

Day 1 visit

Tests: UCVA, SLE (wound secure, corneal edema, AC cell and depth, IOL position, other notable findings), IOP.

Week 1 visit

Tests: UCVA, include UCIVA and UCNVA, MRx, SLE (see Day 1 note above), IOP, DFE.

If contralateral/second eye also has cataract, please fax the following data with your 1-week report to our surgery coordinators:

- Glare data on second eye
- Post-op manifest refraction of first surgical eye

We consider your 1-week findings in planning the second eye surgery, and your prompt response is appreciated.

Once both eyes have been completed measure binocular and monocular UCVA.

Month 1

Tests: UCVA, include UCIVA and UCNVA, MRx, SLE, IOP.

Once both eyes have been completed measure binocular and monocular UCVA.

If vision outcome is not as expected when the patient finishes post-operative drops, please alert NWES and return the patient for evaluation.

Month 4

Tests: UCVA, include UCIVA and UCNVA, MRx, SLE, IOP.

Once both eyes have been completed measure binocular and monocular UCVA. In addition, please indicate the patient's satisfaction with the surgical outcome on your post-operative records.

At 12 months

We recommend a comprehensive exam and ask that you please fax the results of UCVA, UCIVA, UCNVA, and MRX to our clinic.

Please fax your completed exam notes to Northwest Eye Surgeons.